

Lecture 4  
*Simulation of differential-algebraic equations*

Erik Frisk  
erik.frisk@liu.se

Department of Electrical Engineering  
Linköping University

November 23, 2017



*Outline*

---

- *Consistent initial conditions*
- *Structural index*
- *Index reduction with dummy derivatives*
- *Summary and exercises*

*Outline*

---

- *Consistent initial conditions*
- *Structural index*
- *Index reduction with dummy derivatives*
- *Summary and exercises*

*Now, what was the problem with initial conditions?*

---

For an initial value problem for an ODE

$$\dot{x} = f(x, t), \quad x(t_0) = x_0$$

there are no limitations (except domain for  $f$ ) for the initial condition  $x_0$ .

For a DAE

$$F(t, y, \dot{y}) = 0$$

it is not sufficient that  $\dot{y}(t_0)$  and  $y(t_0)$  fulfills

$$F(t_0, y_0, \dot{y}(t_0)) = 0$$

## Now, what was the problem with initial conditions?

For example, remember the DAE from the first DAE lecture

$$\begin{aligned} \dot{x}_1 + x_2 + x_3 &= f_1 & x_1(t) &= f_2(t) - \dot{f}_3(t) \\ \dot{x}_2 + x_1 &= f_2 & x_2(t) &= f_3(t) \\ x_2 &= f_3 & x_3(t) &= f_1(t) - f_3(t) - \dot{f}_2(t) + \ddot{f}_3(t) \end{aligned}$$

Here, no freedom at all and the initial conditions has to satisfy

$$\begin{aligned} x_1(t_0) &= f_2(t_0) - \dot{f}_3(t_0) \\ x_2(t_0) &= f_3(t_0) \\ x_3(t_0) &= f_1(t_0) - f_3(t_0) - \dot{f}_2(t_0) + \ddot{f}_3(t_0) \end{aligned}$$

### Problem

We do not want to solve the DAE to find initial conditions!

## Pantelides algorithm

We know that given a DAE

$$F(\dot{y}, y, t) = 0$$

we can differentiate well chosen equation a suitable number of times to obtain a model including all constraints for the initial condition.

$$\begin{aligned} F(\dot{y}, y, t) &= 0 \\ \frac{d}{dt} F(\dot{y}, y, t) &= 0 \\ \frac{d^2}{dt^2} F(\dot{y}, y, t) &= 0 \\ &\vdots \\ \frac{d^j}{dt^j} F(\dot{y}, y, t) &= 0 \end{aligned}$$

Two questions:

- which equations?
- differentiate how many times?

## Pantelides algorithm - consistent initial conditions

Finding a consistent initial condition  $(y(t_0), \dot{y}(t_0), t_0)$  for a DAE

$$F(y, \dot{y}, t) = 0$$

with unknown index is a difficult problem in general. By differentiation we can obtain “hidden” conditions on the initial condition.

### Pantelides algorithm

Graph theoretical approach to find the conditions that has to be satisfied and solved by a numerical equation solver.

- Good because based on equation structure only, possible to make automatic
- Bad based on equation structure only, does not give analytical results
  
- Can be used to compute differential index
- Can be used for index reduction. Will come back to this.

## The small example again

$$\begin{aligned} e_1 : \dot{x}_1 + x_2 + x_3 &= f_1 \\ e_2 : \dot{x}_2 + x_1 &= f_2 \\ e_3 : x_2 &= f_3 \end{aligned}$$

Differentiate  $e_2$  once and  $e_3$  twice and collect the equations. These 6 equations can be solved for the 6 variables  $(x_1(0), \dot{x}_1(0), x_2(0), \dot{x}_2(0), \ddot{x}_2(0), x_3(0))$  for a consistent initial condition

$$\begin{aligned} e_1 : \dot{x}_1 + x_2 + x_3 &= f_1 \\ e_2 : \dot{x}_2 + x_1 &= f_2 \\ \dot{e}_2 : \ddot{x}_2 + \dot{x}_1 &= \dot{f}_2 \\ e_3 : x_2 &= f_3 \\ \dot{e}_3 : \dot{x}_2 &= \dot{f}_3 \\ \ddot{e}_3 : \ddot{x}_2 &= \ddot{f}_3 \end{aligned}$$

The new DAE  $(e_1, \dot{e}_2, \ddot{e}_3)$  is index 1 (see next slide) and we had to differentiate  $e_3$  twice. This is no coincidence.

## The small example again

---

$$e_1 : \dot{x}_1 + x_2 + x_3 = f_1$$

$$e_2 : \ddot{x}_2 + \dot{x}_1 = \dot{f}_2$$

$$e_3 : \ddot{x}_2 = \ddot{f}_3$$

$\Rightarrow$

$$e_3 : \ddot{x}_2 = \ddot{f}_3$$

$$e_2 : \dot{x}_1 = \dot{f}_2 - \ddot{f}_3$$

$$e_1 : x_3 = f_1 - \dot{x}_1 + x_2 = f_1 - \dot{f}_2 - \ddot{f}_3 + x_2$$

## Initial condition, example, cont.

---

For the model

$$\dot{x}_1 = x_1 + x_2, \quad 0 = x_1 + 2x_2 + a$$

we can not obtain any new constraints on the initial condition by differentiating the equations.

For every new differentiation, we get a new variable. Differentiation gives

$$\ddot{x}_1 = \dot{x}_1 + \dot{x}_2$$

$$0 = \dot{x}_1 + 2\dot{x}_2 + \dot{a}$$

These can *always* be satisfied by choosing a suitable value for  $\ddot{x}_1(t_0)$  and  $\dot{x}_2(t_0)$ .

From this you can conclude that it is sufficient to solve the original equations for  $(x_1(t_0), \dot{x}_1(t_0), x_2(t_0))$ . (This we already knew since the DAE is index 1)

## Initial condition, example

---

For the model (1-DOF)

$$e_1 : \dot{x}_1 + \dot{x}_2 = a(t), \quad e_2 : x_1 + x_2^2 = b(t)$$

we can differentiate equation  $e_2$  to obtain the constraint

$$\dot{e}_2 : \dot{x}_1 + 2x_2\dot{x}_2 = b'(t)$$

and we are done since  $(\dot{x}_1, \dot{x}_2)$  can be solved for in  $(e_1, \dot{e}_2)$ .

The initial condition is therefore obtained by solving

$$\dot{x}_1(t_0) + \dot{x}_2(t_0) = a(t_0)$$

$$x_1(t_0) + x_2^2(t_0) = b(t_0)$$

$$\dot{x}_1(t_0) + 2x_2(t_0)\dot{x}_2(t_0) = \dot{b}(t_0)$$

for  $(x_1(t_0), x_2(t_0), \dot{x}_1(t_0), \dot{x}_2(t_0))$ . With 4 unknowns and three equations, 1-DOF which we kind of knew.

## New constraints, when?

---

What does the situation look like when we can, and cannot, obtain new constraints by differentiating? What separates the two examples?

$$\dot{x}_1 + \dot{x}_2 = a(t)$$

$$x_1 + x_2^2 = b(t)$$

$$\dot{x}_1 = x_1 + x_2$$

$$0 = x_1 + 2x_2 + a$$

New constraints exists

No new constraints

- Difference is the index of the DAE:s
- A **sufficient** condition for a DAE to have at most index 1 is that it is possible to solve for highest differentiated variables.
- For semi-explicit DAE:s it is also a necessary condition

## Differentiation a set of equations

---

Assume a DAE

$$f(x, \dot{x}, y, t) = 0, \quad f : \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^m \times \mathbb{R} \rightarrow \mathbb{R}^{n+m}$$

The highest differentiated variables are interesting, let  $z = (\dot{x}, y)$  be a vector with the highest derivatives

$$f(x, z, t) = 0$$

Find a **subset** with  $k$  equations

$$\bar{f}(\bar{x}, \bar{z}, t) = 0, \quad \bar{x} \in \mathbb{R}^q, \quad \bar{z} \in \mathbb{R}^l$$

Assume a well formed model, e.g., no dependent set of equations.

## Differentiation a set of equations, cont.

---

Since  $r$  (number of new highest derivatives) can not be larger than  $l$  (number of highest derivatives in  $\bar{f}$ ), a sufficient condition for  $k - r > 0$  is

$$l < k$$

The above property implies that the set of equations  $\bar{f}$  contains fewer highest ordered derivatives than equations. Minimally structurally singular (MSS).

Set of equations is **overdetermined** with respect to the highest ordered differentiated variables.

## Differentiation a set of equations, cont.

---

Differentiate the set of equations  $\bar{f}$  w.r.t.  $t$

$$\bar{f}_{\bar{x}} \dot{\bar{x}} + \bar{f}_{\bar{z}} \dot{\bar{z}} + \bar{f}_t = 0$$

The number of “new” highest derivatives  $\bar{z}'$  appearing in the differentiated equation is determined by  $\text{rang } \bar{f}_{\bar{z}}$ .

$$r \leq \min(k, l) = \min(\text{equations, number } z \text{ i } \bar{f})$$

With no dependencies it holds that

$$\text{rang } (\bar{f}_{\bar{x}} \bar{f}_{\bar{z}}) = k$$

The conclusion so far is then:

- we get  $k - r$  new equations/constraints when differentiating  $\bar{f}$ .
- all subsets of equations where  $k - r > 0$  are useful to obtain new constraints

## New constraints, when? Revisited!

---

What does the situation look like when we can, and cannot, obtain new constraints by differentiating? What separates the two examples?

$$e_1 : \dot{x}_1 + \dot{x}_2 = a(t)$$

$$e_2 : x_1 + x_2^2 = b(t)$$

$$e_1 : \dot{x}_1 = x_1 + x_2$$

$$e_2 : 0 = x_1 + 2x_2 + a$$

New constraints exists, equation  $e_2$  contains none of the highest ordered differentiated variables  $\dot{x}_1$  or  $\dot{x}_2$ .

With  $\bar{f}$  equal to  $e_2$  then  $k = 1$  and  $l = r = 0 \Rightarrow$   
 $k - r = 1 - 0 = 1$  new constraints.

No new constraints. Both  $e_1$  and  $e_2$  each contain one of the highest ordered differentiated variables  $\dot{x}_1$  and  $x_2$  respectively.

With  $\bar{f}$  equal to  $e_1$  or  $e_2$  then  $k = 1$  and  $l = r = 1 \Rightarrow$   
 $k - r = 1 - 1 = 0$  new constraints.

## Sketch of Pantelides algorithm

Sketch of the basic principles for Pantelides algorithm for a DAE

$$f(\dot{x}, x, y, t) = 0.$$

- 1 Define  $z = (\dot{x}, y)$
- 2 Find all subsets of equations where  $l < k$ , i.e., overdetermined w.r.t. the highest ordered derivatives. If none exists, exit.
- 3 Differentiate these equations and extend the model with the new equations. Go to step 1.
  - There are many, very many, possible subsets. This has to be done in a smart way to not run into complexity problems.
  - Solvable!

### Pantelides algorithm

A graph theoretical algorithm that do the above efficiently for large systems with no symbolic computations.

An implementation of the algorithm, courtesy Mattias Krysander, can be downloaded from the course website, can come in handy when solving some of the exercises.

## Matching and structural rank

A matrix is said to have full structural rank if all variables can be matched.

	$x_1$	$x_2$	$x_3$
$e_1$	X	X	X
$e_2$	X	X	
$e_3$		X	X

	$x_1$	$x_2$	$x_3$
$e_1$	X	X	X
$e_2$		X	
$e_3$		X	

- If there exists a complete matching with respect to  $x$  in the structure for a function  $f(x, y)$ , the Jacobian  $f_x(x, y)$  has full rank for almost all  $f$  with the same structure.

## Matching - a useful graph theoretical concept

	$x_1$	$x_2$	$x_3$
$f_1(x_1, x_2, x_3, y) = 0$	X	X	X
$f_2(x_1, x_2, y) = 0$	X	X	
$f_3(x_2, x_3, y) = 0$		X	X

Implicit function theorem gives that there exists a local solution for  $x$  in the equation  $f(x, y) = c$  at  $x = x_0$  if

$$\left. \frac{\partial f}{\partial x} \right|_{x=x_0} = \begin{pmatrix} * & * & * \\ * & * & 0 \\ 0 & * & * \end{pmatrix} \Bigg|_{x=x_0}$$

has full column rank.

### Matching

(here) A pairing of equations with variables

## Example: Matching condition for semi-explicit index 1

In a semi-explicit DAE

$$\dot{x} = f(x, y)$$

$$0 = g(x, y)$$

the highest differentiated variables  $z = (\dot{x}, y)$ . DAE has (local) index 0/1 if

$$\left. \frac{\partial}{\partial z} \begin{pmatrix} f(x, y) \\ g(x, y) \end{pmatrix} \right|_{z=z_0}$$

has full column rank.

Convince yourselves that the DAE has structural index 0/1 if the structure of the DAE has a complete matching with respect to the variables  $z$ .

## Pantelides on an index 2 DAE

### Step 1

$$\begin{aligned} e_1 : \dot{x} &= f(x, y) \\ e_2 : \dot{y} &= g(x, y, z) \\ e_3 : 0 &= h(x) \end{aligned}$$

	$\dot{x}$	$\dot{y}$	$z$
$e_1$	X		
$e_2$		X	X
$e_3$			

Highest differentiated variables:  $(\dot{x}, \dot{y}, z)$

### Step 2

$$\begin{aligned} e_1 : \dot{x} &= f(x, y) \\ e_2 : \dot{y} &= g(x, y, z) \\ \dot{e}_3 : 0 &= \frac{d}{dt} h(x) \end{aligned}$$

	$\dot{x}$	$\dot{y}$	$z$
$e_1$	X		
$e_2$		X	X
$\dot{e}_3$	X		

Highest differentiated variables:  $(\dot{x}, \dot{y}, z)$

## Pantelides on an index 2 DAE, cont.

### Step 3

$$\begin{aligned} \dot{e}_1 : \ddot{x} &= \frac{d}{dt} f(x, y) \\ e_2 : \dot{y} &= g(x, y, z) \\ \ddot{e}_3 : 0 &= \frac{d^2}{dt^2} h(x) \end{aligned}$$

	$\ddot{x}$	$\dot{y}$	$z$
$\dot{e}_1$	X	X	
$e_2$		X	X
$\ddot{e}_3$	X		

Highest differentiated vars:  $(\ddot{x}, \dot{y}, z)$

### Resulting system of equations (6 equations in 6 unknowns)

$$\begin{aligned} e_1 : \dot{x} &= f(x, y) \\ \dot{e}_1 : \ddot{x} &= d/dt f(x, y) \\ e_2 : \dot{y} &= g(x, y, z) \\ e_3 : 0 &= h(x) \\ \dot{e}_3 : 0 &= d/dt h(x) \\ \ddot{e}_3 : 0 &= d^2/dt^2 h(x) \end{aligned}$$

	$x$	$\dot{x}$	$\ddot{x}$	$y$	$\dot{y}$	$z$
$e_1$	X	X		X		
$\dot{e}_1$	X	X	X	X	X	
$e_2$	X			X	X	X
$e_3$	X					
$\dot{e}_3$	X	X				
$\ddot{e}_3$	X	X	X			

## Example: Pendulum equations

$$\begin{aligned} \ddot{x} &= Tx \\ \ddot{y} &= Ty - g \\ 0 &= x^2 + y^2 - L^2 \end{aligned} \quad \Rightarrow \quad \begin{aligned} \dot{x} &= w \\ \dot{y} &= z \\ \dot{w} &= Tx \\ \dot{z} &= Ty - g \\ 0 &= x^2 + y^2 - L^2 \end{aligned}$$

Highest differentiated variables are  $(\dot{x}, \dot{y}, \dot{w}, \dot{z}, T)$

	$T$	$\dot{x}$	$\dot{w}$	$\dot{y}$	$\dot{z}$
$e_1$		X			
$e_2$				X	
$e_3$	X		X		
$e_4$	X				X
$e_5$					

## Differentiate $e_5$ and extend the model

$$\begin{aligned} e_1 : \dot{x} &= w & e'_5 : 0 &= 2x\dot{x} + 2y\dot{y} \\ e_2 : \dot{y} &= z \\ e_3 : \dot{w} &= Tx \\ e_4 : \dot{z} &= Ty - g \\ e_5 : 0 &= x^2 + y^2 - L^2 \end{aligned}$$

Highest differentiated variables are  $(\dot{x}, \dot{y}, \dot{w}, \dot{z}, T)$

	$T$	$\dot{x}$	$\dot{w}$	$\dot{y}$	$\dot{z}$
$e_1$		X			
$e_2$				X	
$e_3$	X		X		
$e_4$	X				X
$e'_5$		X		X	

## Differentiate $e_1$ , $e_2$ , and $e_5$

$$\begin{array}{ll}
 e_1 : & \dot{x} = w \\
 e_2 : & \dot{y} = z \\
 e_3 : & \dot{w} = Tx \\
 e_4 : & \dot{z} = Ty - g \\
 e_5 : & 0 = x^2 + y^2 - L^2 \\
 \ddot{e}_5 : & 0 = 2x\dot{x} + 2y\dot{y} \\
 \ddot{e}_2 : & 0 = 2\dot{x}^2 + 2x\ddot{x} + 2\dot{y}^2 + 2y\ddot{y}
 \end{array}$$

	$T$	$\dot{w}$	$\dot{z}$	$\ddot{x}$	$\ddot{y}$
$e_3$	X	X			
$\dot{e}_1$		X		X	
$e_4$	X		X		
$\ddot{e}_5$				X	X
$\ddot{e}_2$			X		X

$$\nu = (1, 1, 0, 0, 2)$$

11 variables and 9 equations, i.e., 2 degrees of freedom. Makes sense ( $\approx$  position and velocity)

## Structural index

Can be computed by Pantelides algorithm. Determine how in exercise 2.13. (an error in the book by P. Fritzson so you will have to solve it by yourselves)

Remember the non-trivial relationship between index  $\nu$  and structural index  $\nu_{str}$ .

$$\nu = \nu_{str}, \quad \nu < \nu_{str}, \quad \nu > \nu_{str}$$

All is possible!

## Outline

- Consistent initial conditions
- Structural index
- Index reduction with dummy derivatives
- Summary and exercises

## Outline

- Consistent initial conditions
- Structural index
- Index reduction with dummy derivatives
- Summary and exercises

## Index reduction

- Differentiate to the underlying ODE is often a not satisfactory solution
- due to that the underlying ODE has a larger solution set
- requires stabilization/projection techniques to avoid violating algebraic constraints
- Objective is then to do index reduction while keeping solution set
- To save all differentiated equations, and not only the underlying ODE, is one such way. The result is a **overdetermined** index-1 DAE. Consistency etc. is typically violated at discretization, projection etc. is needed.
- Dummy derivatives is a method that solves such problems.

## Index reduction with dummy derivatives, principle

$$\begin{array}{l}
 F(\dot{x}, x, t) = 0 \\
 \vdots \\
 \frac{d^j}{dt^j} F(\dot{x}, x, t) = 0
 \end{array}$$

If only  $j$  is large enough (index) the equations are an index-1 DAE with the exact same solution set as the original DAE. Problem: system is (violently) overdetermined.

### Principle for index reduction

- 1 Let Pantelides algorithm determine the number of times to differentiate
- 2 Differentiate equations according to Pantelides, collect all equations
- 3 Simplified: For each differentiated equation, introduce an algebraic variable such that the system becomes exactly determined
- 4 Result: exactly determined index 1 DAE with the same solution set as the original DAE

## Small example that shows the principle

Index-1 system

$$\begin{array}{l}
 e_1 : \dot{x}_1 + \dot{x}_2 = a(t) \\
 e_2 : x_1 + x_2^2 = b(t)
 \end{array}$$

Differentiate  $e_2$  once gives the *overdetermined* system of equations

$$\begin{array}{l}
 e_1 : \dot{x}_1 + \dot{x}_2 = a(t) \\
 e_2 : x_1 + x_2^2 = b(t) \\
 \dot{e}_2 : \dot{x}_1 + 2x_2 \dot{x}_2 = \dot{b}(t)
 \end{array}$$

Replace  $\dot{x}_1$  for a *new algebraic* variable  $x'_1$

$$\begin{array}{l}
 e_1 : x'_1 + \dot{x}_2 = a(t) \\
 e_2 : x_1 + x_2^2 = b(t) \\
 \dot{e}_2 : x'_1 + 2x_2 \dot{x}_2 = \dot{b}(t)
 \end{array}
 \Rightarrow
 \begin{array}{l}
 x_1 = b(t) - x_2^2 \\
 \dot{x}_2 = \frac{\dot{b}(t) - a(t)}{2x_2 - 1}
 \end{array}$$

and solve for  $(x_1, x'_1, x_2)$ . Can be proven to have the *same* solution set as the original equations.

## Example - a DAE with index 3

$$\begin{array}{l}
 (a) : \dot{x} = y \\
 (b) : \dot{y} = z \\
 (c) : x = f(t)
 \end{array}
 \qquad
 \begin{array}{l}
 x(t) = f(t) \\
 y(t) = \dot{f}(t) \\
 z(t) = \ddot{f}(t)
 \end{array}$$

Pantelides states that we should differentiate (c) twice and (a) once. Collecting the equations

$$\begin{array}{l}
 (c) : x = f(t) \\
 (\dot{c}) : \dot{x} = \dot{f}(t) \\
 (\ddot{c}) : \ddot{x} = \ddot{f}(t) \\
 (a) : \dot{x} = y \\
 (\dot{a}) : \ddot{x} = \dot{y} \\
 (b) : \dot{y} = z
 \end{array}$$



### Example, cont.

The differentiated model is overdetermined (3 unknowns, 6 equations).  
Introduce an algebraic variable for each differentiated equation

$$x' = \dot{x}, x'' = \ddot{x}, y' = \dot{y}$$

**Important!!** Variables  $x'$ ,  $x''$ ,  $y'$  is here **algebraic** variables.

(c) :	$x = f(t)$
( $\dot{c}$ ) :	$x' = \dot{f}(t)$
( $\ddot{c}$ ) :	$x'' = \ddot{f}(t)$
(a) :	$x' = y$
( $\dot{a}$ ) :	$x'' = y'$
(b) :	$y' = z$

Exactly determined, index 1 DAE (6 unknowns, 6 equations)

Somewhat extreme example where the system turns into a purely algebraic system of equations, but it illustrates a simple case.

### Structure of the differentiated system

Simple if

- Pantelides algorithm only differentiates 1 time and only one new variable is introduced
- How should the situation where equations are differentiated more than once and multiple new variables are introduced simultaneously?


### Principle for index reduction

- 1 Let Pantelides algorithm determine the number of times to differentiate
- 2 Differentiate equations according to Pantelides, collect all equations
- 3 Simplified: For each differentiated equation, introduce an algebraic variable such that the system becomes exactly determined
- 4 Result: exactly determined index 1 DAE with the same solution set as the original DAE

Step 3 need to be clarified.

### Structure of the differentiated system

take the differentiated system, by permutations of equations and variables you can always get a Block Lower Triangle (BLT) form w.r.t. the most differentiated variables



Consider one block at a time.

### Example - a system of index 2

---

$$\begin{aligned}(a) : & \quad x_1 + x_2 + u_1 = 0 \\(b) : & \quad x_1 + x_2 + x_3 + u_2 = 0 \\(c) : & \quad x_1 + \dot{x}_3 + x_4 + u_3 = 0 \\(d) : & \quad 2\ddot{x}_1 + \ddot{x}_2 + \ddot{x}_3 + \dot{x}_4 + u_4 = 0\end{aligned}$$

Pantelides gives  $\nu = (2, 2, 1, 0)$  and the differentiated system  $\mathcal{G}x = 0$  is:

$$\begin{aligned}(\ddot{a}) : & \quad \ddot{x}_1 + \ddot{x}_2 + \ddot{u}_1 = 0 \\(\ddot{b}) : & \quad \ddot{x}_1 + \ddot{x}_2 + \ddot{x}_3 + \ddot{u}_2 = 0 \\(\ddot{c}) : & \quad \dot{x}_1 + \ddot{x}_3 + \dot{x}_4 + \dot{u}_3 = 0 \\(\ddot{d}) : & \quad 2\ddot{x}_1 + \ddot{x}_2 + \ddot{x}_3 + \dot{x}_4 + u_4 = 0\end{aligned}$$

We have introduced  $2+2+1=5$  equations, i.e., we need to introduce 5 dummy variables. Which ones? Not as easy as in the first example where there was a one-to-one relation between differentiated equation and new variable. Candidates are  $(\dot{x}_1, \ddot{x}_1, \dot{x}_2, \ddot{x}_2, \dot{x}_3, \ddot{x}_3, \dot{x}_4)$ , which 5 to choose?

### Example - a system of index 2, cont.

---

We can choose  $(\ddot{x}_1, \ddot{x}_3, \dot{x}_4)$  or  $(\ddot{x}_2, \ddot{x}_3, \dot{x}_4)$ . Choose

$$\hat{z}^{[1]} = (\ddot{x}_1, \ddot{x}_3, \dot{x}_4)$$

these variables will be introduced as dummy variables in the final DAE. We are not done since we yet only have define 3 dummy variables, we must have 5.

Now look at the differentiated equations, with one less differentiation (these also are part of the system)

$$\begin{aligned}(\dot{a}) : & \quad \dot{x}_1 + \dot{x}_2 + \dot{u}_1 = 0 \\(\dot{b}) : & \quad \dot{x}_1 + \dot{x}_2 + \dot{x}_3 + \dot{u}_2 = 0 \\(\dot{c}) : & \quad x_1 + \dot{x}_3 + x_4 + u_3 = 0\end{aligned}$$

Candidates for new dummy variables are  $(\dot{x}_1, \dot{x}_3, x_4)$ . Analyze this sub-model in the same way as before.

### Example - a system of index 2, cont.

---

The differentiated system  $\mathcal{G}x = 0$  is **not** of the type with one-to-one relation between differentiated equation and variable.

The highest differentiated variables are  $(\ddot{x}_1, \ddot{x}_2, \ddot{x}_3, \dot{x}_4)$  and  $\mathcal{G}$  consists of a block  $g_1$ , w.r.t. the highest differentiated variables  $z_1$

$$\frac{\partial g_1}{\partial z_1} = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \\ 2 & 1 & 1 & 1 \end{pmatrix}$$

In these, the three first equations are differentiated; get that part

$$H_1 = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}$$

Choice of variables should be such that index  $\leq 1$  is retained, i.e., all highest differentiated variables should still be matched. We must choose variables such that the corresponding sub-matrix of  $H_1$  has full rank.

### Example - a system of index 2, cont.

---

Extract the differentiated equations

$$\begin{aligned}(\dot{a}) : & \quad \dot{x}_1 + \dot{x}_2 + \dot{u}_1 = 0 \\(\dot{b}) : & \quad \dot{x}_1 + \dot{x}_2 + \dot{x}_3 + \dot{u}_2 = 0\end{aligned}$$

Highest derivatives are  $z_1 = (\dot{x}_1, \dot{x}_3, x_4)$ . Differentiate w.r.t.  $z_1$  to obtain

$$H_1^{[2]} = \begin{pmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \end{pmatrix}$$

and here we must choose

$$\hat{z}_1^{[2]} = (\dot{x}_1, \dot{x}_3)$$

as dummy variables and then we have selected our 5 and are done!

### Example - a system of index 2, cont.

---

The final index 1 DAE is then

$$\begin{aligned}(a) : & \quad x_1 + x_2 + u_1 = 0 \\ (\dot{a}) : & \quad x_1' + \dot{x}_2 + \dot{u}_1 = 0 \\ (\ddot{a}) : & \quad x_1'' + \ddot{x}_2 + \ddot{u}_1 = 0 \\ (b) : & \quad x_1 + x_2 + x_3 + u_2 = 0 \\ (\dot{b}) : & \quad x_1' + \dot{x}_2 + x_3' + \dot{u}_2 = 0 \\ (\ddot{b}) : & \quad x_1'' + \ddot{x}_2 + x_3'' + \ddot{u}_2 = 0 \\ (c) : & \quad x_1 + x_3' + x_4 + u_3 = 0 \\ (\dot{c}) : & \quad x_1' + x_3'' + x_4' + \dot{u}_3 = 0 \\ (d) : & \quad 2x_1'' + \ddot{x}_2 + x_3'' + x_4' + u_4 = 0\end{aligned}$$

which has the same solution set as the original index 2 DAE.

The 9 unknown variables are  $(x_1, x_1', x_1'', x_2, x_3, x_3', x_3'', x_4, x_4')$  out of which all are algebraic except  $x_2$  which appears differentiated of order 2.

### Dummy derivatives summary

---

- Pantelides algorithm plus a procedure to introduce algebraic variables gives a low-index system with the same solution set as the original DAE
- Is all index related issues thereby solved?
- Pros/cons
- Structural and analytical steps

### Example - a system of index 2, cont.

---

It is possible to get a simpler solution. Pantelides differentiated for an ODE, it is really sufficient to differentiate until an index 1 DAE.

Rather straightforward changes to the basic principle gives the somewhat simpler system

$$\begin{aligned}(a) : & \quad x_1 + x_2 + u_1 = 0 \\ (\dot{a}) : & \quad x_1' + \dot{x}_2 + \dot{u}_1 = 0 \\ (b) : & \quad x_1 + x_2 + x_3 + u_2 = 0 \\ (\dot{b}) : & \quad x_1' + \dot{x}_2 + x_3' + \dot{u}_2 = 0 \\ (c) : & \quad x_1 + x_3' + x_4 + u_3 = 0 \\ (d) : & \quad 2x_1' + \ddot{x}_2 + \dot{x}_3 + \dot{x}_4 + u_4 = 0\end{aligned}$$

### Outline

---

- Consistent initial conditions
- Structural index
- Index reduction with dummy derivatives
- Summary and exercises

- Three problems have been discussed:
  - consistent initial conditions
  - determining index
  - index reduction
- Pantelides algorithm: a graph theoretical algorithm to find the system of equations to solve for consistent initial conditions given a DAE of arbitrary index
- Pantelides algorithm is a cornerstone for solving all three problems
- Well suited for implementation in a general purpose DAE simulator
- Structural results, not analytical!

Mandatory exercises (marked in the exercises): 2.13, 2.29

### *Lecture 4*

## *Simulation of differential-algebraic equations*

Erik Frisk

`erik.frisk@liu.se`

Department of Electrical Engineering  
Linköping University

November 23, 2017